

DAA2 Human, Gastrointestinal stromal tumor, primary-metastasis-normal

No.	Age	Sex	Organ	Diagnosis	pTNM	Stage	Months*	Tissue type**
1	45	F	Jejunum	GIST, high risk	T3N0M0	III B	.	6
2	57	M	Jejunum	GIST, high risk	T4N0M0	III B	.	6
3	77	F	Stomach	GIST, high risk	T4N0M0	III B	.	6
4	60	F	Stomach	GIST, high risk	T3N0M0	III A	.	6
5	59	M	Jejunum	GIST, high risk	T3N0M0	III B	.	6
6	58	M	Small intestine	GIST, high risk	T4N1M0	IV	.	6
7	65	F	Stomach	GIST, low risk	T3N0M0	I B	.	4
8	61	M	Stomach	GIST, high risk	T4N1M0	IV	.	6
9	65	F	Stomach	GIST, high risk	T4N0M0	II	.	6
10	44	M	Small intestine	GIST, high risk	T3N0M0	III B	.	6
11	53	F	Duodenum	GIST, intermediate risk	T3N0M0	II	.	5
12	50	M	Stomach	GIST, low risk	T2N0M0	I A	.	4
13	68	M	Stomach	GIST, high risk	T4N0M0	III B	.	6
14	34	M	Duodenum	GIST, high risk	T2N0M0	II	.	6
15	61	M	Stomach	GIST, intermediate risk	T3N0M0	I B	.	5
16	58	F	Stomach	GIST, high risk	T3N0M0	III A	.	6
17	57	F	Stomach	GIST, low risk	T2N0M0	I A	.	4
18	40	M	Stomach	GIST, high risk	T4N0M0	III B	.	6
19	66	F	Rectum	GIST, high risk	T3N0M0	III B	.	6
20	45	F	Abdominal cavity	GIST, high risk	T4N0M1	IV	.	6
21	66	M	Stomach	GIST, high risk	T4N0M0	III B	.	6
22	83	F	Stomach	GIST, high risk	T4N0M0	III B	.	6
23	60	F	Jejunum	GIST, intermediate risk	T3N0M0	II	.	5
24	65	M	Jejunum	GIST, high risk	T4N0M0	III B	.	6
25	56	F	Duodenum	GIST, low risk	T2N0M0	I	.	4
26	85	F	Stomach	GIST, high risk	T4N0M0	II	.	6
27	67	F	Duodenum	GIST, intermediate risk	T3N0M0	II	.	5
28	52	M	Stomach	GIST, high risk	T4N0M0	III B	.	6
29	65	M	Stomach	GIST, low risk	T2N0M0	I A	.	4
30	65	M	Ileum	GIST, intermediate risk	T3N0M0	II	.	5
31	45	F	Stomach	GIST, low risk	T2N0M0	I A	.	4
32	55	M	Rectum	GIST, high risk	T3N0M0	III B	.	6
33	71	M	Stomach	GIST, high risk	T4N0M0	II	.	6
34	67	F	Stomach	GIST, intermediate risk	T2N0M0	II	.	5
35	62	F	Stomach	GIST, intermediate risk	T3N0M0	I B	.	5
36	48	M	Stomach	GIST, high risk	T4N0M0	III B	.	6
37	80	M	Small intestine	GIST, low risk	T2N0M0	I	.	4
38	60	M	Small intestine	GIST, intermediate risk	T3N0M0	II	.	5
39	43	M	Duodenum	GIST, low risk	T3N0M0	II	.	4
40	53	F	Stomach	GIST, high risk	T3N0M0	III A	.	6
41	78	M	Liver	spindle cell sarcoma, metastatic	M1	IV	0	6
42	81	F	Lung	leiomyosarcoma, metastatic	M1	IV	0	6
43	65	M	Omentum	GIST, recurred	T4N0M0	III B	75	6
44	80	M	Liver	GIST, metastatic	M1	IV	48	6
45	64	M	Mesentery	GIST, metastatic	T4N0M0	III B	28	6
46	67	M	Mesentery	GIST, metastatic	T2N0M0	I	48	6
47	35	M	Ileum	GIST, recurred	T3N0M0	III B	36	6
48	62	M	Omentum	GIST, metastatic	T4N0M0	III B	0	6
49	45	F	Liver	GIST, metastatic (match of #1)	M1	IV	80	6
50	57	M	Mesentery	GIST, low risk, recurred (match of #2)	T2N0M0	I	16	4
51	77	F	Stomach	normal (match of #3)	.	.	.	3
52	60	F	Stomach	normal (match of #4)	.	.	.	3
53	59	M	Jejunum	normal (match of #5)	.	.	.	3
54	58	M	Small intestine	normal (match of #6)	.	.	.	3
55	61	M	Stomach	normal (match of #8)	.	.	.	3
56	44	M	Small intestine	normal (match of #10)	.	.	.	3
57	68	M	Stomach	normal (match of #13)	.	.	.	3
58	66	F	Rectum	normal (match of #19)	.	.	.	3
59	65	M	Jejunum	normal (match of #24)	.	.	.	3
60	.	.	Carbon

TNM and Stage: AJCC Cancer Staging Manual (7th Edition)

Months*: interval between primary and metastatic/recurrent cancer

Tissue type**: see General information sheet

Supplementary data sheet contains tumor size, mitosis number, immunohistochemistry result, and survival