



## ZG2 Normal organs of adult mice, perfusion fixation

No.	Sex	Organ	Abnormal finding
1	M	Skin	.
2	F	Skin	.
3	M	Spleen	.
4	F	Spleen	.
5	M	Skeletal muscle (abdominal wall)	.
6	F	Skeletal muscle (abdominal wall)	.
7	M	Trachea	.
8	F	Trachea	.
9	M	Lung	.
10	F	Lung	.
11	M	Heart	.
12	F	Heart	.
13	M	Salivary gland	.
14	F	Salivary gland	.
15	M	Liver	.
16	F	Liver	.
17	M	Pancreas	.
18	F	Pancreas	.
19	M	Esophagus	.
20	F	Esophagus	.
21	M	Forestomach	.
22	F	Forestomach	.
23	M	Glandular stomach	.
24	F	Glandular stomach	.
25	M	Duodenum	.
26	F	Duodenum	.
27	M	Jejunum	.
28	F	Jejunum	.
29	M	Ileum	.
30	F	Ileum	.
31	M	Colon	.
32	F	Colon	.
33	M	Cecum	.
34	F	Cecum	.
35	M	Rectum	.
36	F	Rectum	.
37	M	Kidney (cortex)	.
38	F	Kidney (cortex)	.
39	M	Kidney (medulla)	.
40	F	Kidney (medulla)	.
41	M	Bladder	.
42	F	Bladder	.
43	M	Prostate	.
44	M	Seminal vesicle	.
45	M	Testis	.
46	F	Uterus	.
47	F	Ovary	.
48	M	Adrenal gland	.
49	F	Adrenal gland	.
50	F	Thymus	.
51	M	Cerebrum	.
52	F	Cerebrum	.
53	M	Cerebellum	.
54	F	Cerebellum	.
55	M	Brain stem	.
56	M	Eyeball	.
57	F	Eyeball	.
58	M	Ear lobe	.
59	F	Ear lobe	.
60	.	Carbon	.

ICR mice, 8 weeks old

Perfusion fixation with 4% paraformaldehyde

Suitable for in situ hybridization and immunohistochemistry